

Mask use in the context of COVID-19

Interim guidance

1 December 2020



This document, which is an update of the guidance published on 5 June 2020, includes new scientific evidence relevant to the use of masks for reducing the spread of SARS-CoV-2, the virus that causes COVID-19, and practical considerations. It contains updated evidence and guidance on the following:

- mask management;
- SARS-CoV-2 transmission;
- masking in health facilities in areas with community, cluster and sporadic transmission;
- mask use by the public in areas with community and cluster transmission;
- alternatives to non-medical masks for the public;
- exhalation valves on respirators and non-medical masks;
- mask use during vigorous intensity physical activity;
- essential parameters to be considered when manufacturing non-medical masks (Annex).

Key points

- The World Health Organization (WHO) advises the use of masks as part of a comprehensive package of prevention and control measures to limit the spread of SARS-CoV-2, the virus that causes COVID-19. A mask alone, even when it is used correctly, is insufficient to provide adequate protection or source control. Other infection prevention and control (IPC) measures include hand hygiene, physical distancing of at least 1 metre, avoidance of touching one's face, respiratory etiquette, adequate ventilation in indoor settings, testing, contact tracing, quarantine and isolation. Together these measures are critical to prevent human-to-human transmission of SARS-CoV-2.
- Depending on the type, masks can be used either for protection of healthy persons or to prevent onward transmission (source control).
- WHO continues to advise that anyone suspected or confirmed of having COVID-19 or awaiting viral laboratory test results should wear a medical mask when in the presence of others (this does not apply to those awaiting a test prior to travel).
- For any mask type, appropriate use, storage and cleaning or disposal are essential to ensure that they are as effective as possible and to avoid an increased transmission risk.

Mask use in health care settings

- WHO continues to recommend that health workers (1) providing care to suspected or confirmed COVID-19

patients wear the following types of mask/respirator in addition to other personal protective equipment that are part of standard, droplet and contact precautions:

- medical mask in the absence of aerosol generating procedures (AGPs)
- respirator, N95 or FFP2 or FFP3 standards, or equivalent in care settings for COVID-19 patients where AGPs are performed; these may be used by health workers when providing care to COVID-19 patients in other settings if they are widely available and if costs is not an issue.
- In areas of known or suspected community or cluster SARS-CoV-2 transmission WHO advises the following:
 - universal masking for all persons (staff, patients, visitors, service providers and others) within the health facility (including primary, secondary and tertiary care levels; outpatient care; and long-term care facilities)
 - wearing of masks by inpatients when physical distancing of at least 1 metre cannot be maintained or when patients are outside of their care areas.
- In areas of known or suspected sporadic SARS-CoV-2 transmission, health workers working in clinical areas where patients are present should continuously wear a medical mask. This is known as targeted continuous medical masking for health workers in clinical areas;
- Exhalation valves on respirators are discouraged as they bypass the filtration function for exhaled air by the wearer.

Mask use in community settings

- Decision makers should apply a risk-based approach when considering the use of masks for the general public.
- In areas of known or suspected community or cluster SARS-CoV-2 transmission:
 - WHO advises that the general public should wear a non-medical mask in indoor (e.g. shops, shared workplaces, schools - see Table 2 for details) or outdoor settings where physical distancing of at least 1 metre cannot be maintained.
 - If indoors, unless ventilation has been assessed to be adequate¹, WHO advises that the general public should wear a non-medical mask, regardless of whether physical distancing of at least 1 metre can be maintained.

¹ For adequate ventilation refer to regional or national institutions or heating, refrigerating and air-conditioning societies enacting ventilation requirements. If not available or applicable, a

recommended ventilation rate of 10 l/s/person should be met (except healthcare facilities which have specific requirements). For more information consult "Coronavirus (COVID-19) response

- Individuals/people with higher risk of severe complications from COVID-19 (individuals $\geq$ 60 years old and those with underlying conditions such as cardiovascular disease or diabetes mellitus, chronic lung disease, cancer, cerebrovascular disease or immunosuppression) should wear medical masks when physical distancing of at least 1 metre cannot be maintained.
- In any transmission scenarios:
 - Caregivers or those sharing living space with people with suspected or confirmed COVID-19, regardless of symptoms, should wear a medical mask when in the same room.

Mask use in children (2)

- Children aged up to five years should not wear masks for source control.
- For children between six and 11 years of age, a risk-based approach should be applied to the decision to use a mask; factors to be considered in the risk-based approach include intensity of SARS-CoV-2 transmission, child's capacity to comply with the appropriate use of masks and availability of appropriate adult supervision, local social and cultural environment, and specific settings such as households with elderly relatives, or schools.
- Mask use in children and adolescents 12 years or older should follow the same principles as for adults.
- Special considerations are required for immunocompromised children or for paediatric patients with cystic fibrosis or certain other diseases (e.g., cancer), as well as for children of any age with developmental disorders, disabilities or other specific health conditions that might interfere with mask wearing.

Manufacturing of non-medical (fabric) masks (Annex)

- Homemade fabric masks of three-layer structure (based on the fabric used) are advised, with each layer providing a function: 1) an innermost layer of a hydrophilic material 2) an outermost layer made of hydrophobic material 3) a middle hydrophobic layer which has been shown to enhance filtration or retain droplets.
- Factory-made fabric masks should meet the minimum thresholds related to three essential parameters: filtration, breathability and fit.
- Exhalation valves are discouraged because they bypass the filtration function of the fabric mask rendering it unserviceable for source control.

Methodology for developing the guidance

Guidance and recommendations included in this document are based on published WHO guidelines (in particular the WHO Guidelines on infection prevention and control of epidemic- and pandemic-prone acute respiratory infections in health care) (2) and ongoing evaluations of all available scientific evidence by the WHO ad hoc COVID-19 Infection Prevention and Control Guidance Development Group (COVID-19 IPC GDG) (see acknowledgement section for list of GDG members). During emergencies WHO publishes interim guidance, the development of which follows a

transparent and robust process of evaluation of the available evidence on benefits and harms. This evidence is evaluated through expedited systematic reviews and expert consensus-building through weekly GDG consultations, facilitated by a methodologist and, when necessary, followed up by surveys. This process also considers, as much as possible, potential resource implications, values and preferences, feasibility, equity, and ethics. Draft guidance documents are reviewed by an external review panel of experts prior to publication.

Purpose of the guidance

This document provides guidance for decision makers, public health and IPC professionals, health care managers and health workers in health care settings (including long-term care and residential), for the public and for manufacturers of non-medical masks (Annex). It will be revised as new evidence emerges.

WHO has also developed comprehensive guidance on IPC strategies for health care settings (3), long-term care facilities (LTCF) (4), and home care (5).

Background

The use of masks is part of a comprehensive package of prevention and control measures that can limit the spread of certain respiratory viral diseases, including COVID-19. Masks can be used for protection of healthy persons (worn to protect oneself when in contact with an infected individual) or for source control (worn by an infected individual to prevent onward transmission) or both.

However, the use of a mask alone, even when correctly used (see below), is insufficient to provide an adequate level of protection for an uninfected individual or prevent onward transmission from an infected individual (source control). Hand hygiene, physical distancing of at least 1 metre, respiratory etiquette, adequate ventilation in indoor settings, testing, contact tracing, quarantine, isolation and other infection prevention and control (IPC) measures are critical to prevent human-to-human transmission of SARS-CoV-2, whether or not masks are used (6).

Mask management

For any type of mask, appropriate use, storage and cleaning, or disposal are essential to ensure that they are as effective as possible and to avoid any increased risk of transmission. Adherence to correct mask management practices varies, reinforcing the need for appropriate messaging (7).

WHO provides the following guidance on the correct use of masks:

- Perform hand hygiene before putting on the mask.
- Inspect the mask for tears or holes, and do not use a damaged mask.
- Place the mask carefully, ensuring it covers the mouth and nose, adjust to the nose bridge and tie it securely to minimize any gaps between the face and the mask. If using ear loops, ensure these do not cross over as this widens the gap between the face and the mask.

- Avoid touching the mask while wearing it. If the mask is accidentally touched, perform hand hygiene.
- Remove the mask using the appropriate technique. Do not touch the front of the mask, but rather untie it from behind.
- Replace the mask as soon as it becomes damp with a new clean, dry mask.
- Either discard the mask or place it in a clean plastic resealable bag where it is kept until it can be washed and cleaned. Do not store the mask around the arm or wrist or pull it down to rest around the chin or neck.
- Perform hand hygiene immediately afterward discarding a mask.
- Do not re-use single-use mask.
- Discard single-use masks after each use and properly dispose of them immediately upon removal.
- Do not remove the mask to speak.
- Do not share your mask with others.
- Wash fabric masks in soap or detergent and preferably hot water (at least 60° Centigrade/140° Fahrenheit) at least once a day. If it is not possible to wash the masks in hot water, then wash the mask in soap/detergent and room temperature water, followed by boiling the mask for 1 minute.

Scientific evidence

Transmission of the SARS-CoV-2 virus

Knowledge about transmission of the SARS-CoV-2 virus is evolving continuously as new evidence accumulates. COVID-19 is primarily a respiratory disease, and the clinical spectrum can range from no symptoms to severe acute respiratory illness, sepsis with organ dysfunction and death.

According to available evidence, SARS-CoV-2 mainly spreads between people when an infected person is in close contact with another person. Transmissibility of the virus depends on the amount of viable virus being shed and expelled by a person, the type of contact they have with others, the setting and what IPC measures are in place. The virus can spread from an infected person's mouth or nose in small liquid particles when the person coughs, sneezes, sings, breathes heavily or talks. These liquid particles are different sizes, ranging from larger 'respiratory droplets' to smaller 'aerosols.' Close-range contact (typically within 1 metre) can result in inhalation of, or inoculation with, the virus through the mouth, nose or eyes (8-13).

There is limited evidence of transmission through fomites (objects or materials that may be contaminated with viable virus, such as utensils and furniture or in health care settings a stethoscope or thermometer) in the immediate environment around the infected person (14-17). Nonetheless, fomite transmission is considered a possible mode of transmission for SARS-CoV-2, given consistent finding of environmental contamination in the vicinity of people infected with SARS-CoV-2 and the fact that other coronaviruses and respiratory viruses can be transmitted this way (12).

Aerosol transmission can occur in specific situations in which procedures that generate aerosols are performed. The scientific community has been actively researching whether the SARS-CoV-2 virus might also spread through aerosol transmission in the absence of aerosol generating procedures (AGPs) (18, 19). Some studies that performed air sampling in

clinical settings where AGPs were not performed found virus RNA, but others did not. The presence of viral RNA is not the same as replication- and infection-competent (viable) virus that could be transmissible and capable of sufficient inoculum to initiate invasive infection. A limited number of studies have isolated viable SARS-CoV-2 from air samples in the vicinity of COVID-19 patients (20, 21).

Outside of medical facilities, in addition to droplet and fomite transmission, aerosol transmission can occur in specific settings and circumstances, particularly in indoor, crowded and inadequately ventilated spaces, where infected persons spend long periods of time with others. Studies have suggested these can include restaurants, choir practices, fitness classes, nightclubs, offices and places of worship (12).

High quality research is required to address the knowledge gaps related to modes of transmission, infectious dose and settings in which transmission can be amplified. Currently, studies are underway to better understand the conditions in which aerosol transmission or superspreading events may occur.

Current evidence suggests that people infected with SARS-CoV-2 can transmit the virus whether they have symptoms or not. However, data from viral shedding studies suggest that infected individuals have highest viral loads just before or around the time they develop symptoms and during the first 5-7 days of illness (12). Among symptomatic patients, the duration of infectious virus shedding has been estimated at 8 days from the onset of symptoms (22-24) for patients with mild disease, and longer for severely ill patients (12). The period of infectiousness is shorter than the duration of detectable RNA shedding, which can last many weeks (17).

The incubation period for COVID-19, which is the time between exposure to the virus and symptom onset, is on average 5-6 days, but can be as long as 14 days (25, 26).

Pre-symptomatic transmission – from people who are infected and shedding virus but have not yet developed symptoms – can occur. Available data suggest that some people who have been exposed to the virus can test positive for SARS-CoV-2 via polymerase chain reaction (PCR) testing 1-3 days before they develop symptoms (27). People who develop symptoms appear to have high viral loads on or just prior to the day of symptom onset, relative to later on in their infection (28).

Asymptomatic transmission – transmission from people infected with SARS-CoV-2 who never develop symptoms – can occur. One systematic review of 79 studies found that 20% (17–25%) of people remained asymptomatic throughout the course of infection. (28). Another systematic review, which included 13 studies considered to be at low risk of bias, estimated that 17% of cases remain asymptomatic (14%–20%) (30). Viable virus has been isolated from specimens of pre-symptomatic and asymptomatic individuals, suggesting that people who do not have symptoms may be able to transmit the virus to others. (25, 29-37)

Studies suggest that asymptomatically infected individuals are less likely to transmit the virus than those who develop symptoms (29). A systematic review concluded that individuals who are asymptomatic are responsible for transmitting fewer infections than symptomatic and pre-symptomatic cases (38). One meta-analysis estimated that there is a 42% lower relative risk of asymptomatic transmission compared to symptomatic transmission (30).

Guidance on mask use in health care settings

Masks for use in health care settings

Medical masks are defined as surgical or procedure masks that are flat or pleated. They are affixed to the head with straps that go around the ears or head or both. Their performance characteristics are tested according to a set of standardized test methods (ASTM F2100, EN 14683, or equivalent) that aim to balance high filtration, adequate breathability and optionally, fluid penetration resistance (39, 40).

Filtering facepiece respirators (FFR), or respirators, offer a balance of filtration and breathability. However, whereas medical masks filter 3 micrometre droplets, respirators must filter more challenging 0.075 micrometre solid particles. European FFRs, according to standard EN 149, at FFP2 performance there is filtration of at least 94% solid NaCl particles and oil droplets. US N95 FFRs, according to NIOSH 42 CFR Part 84, filter at least 95% NaCl particles. Certified FFRs must also ensure unhindered breathing with maximum resistance during inhalation and exhalation. Another important difference between FFRs and other masks is the way filtration is tested. Medical mask filtration tests are performed on a cross-section of the masks, whereas FFRs are tested for filtration across the entire surface. Therefore, the layers of the filtration material and the FFR shape, which ensure the outer edges of the FFR seal around wearer's face, result in guaranteed filtration as claimed. Medical masks, by contrast, have an open shape and potentially leaking structure. Other FFR performance requirements include being within specified parameters for maximum CO₂ build up, total inward leakage and tensile strength of straps (41, 42).

A. Guidance on the use of medical masks and respirators to provide care to suspected or confirmed COVID-19 cases

Evidence on the use of mask in health care settings

Systematic reviews have reported that the use of N95/P2 respirators compared with the use of medical masks (see mask definitions, above) is not associated with statistically significant differences for the outcomes of health workers acquiring clinical respiratory illness, influenza-like illness (risk ratio 0.83, 95%CI 0.63-1.08) or laboratory-confirmed influenza (risk ratio 1.02, 95%CI 0.73-1.43); harms were poorly reported and limited to discomfort associated with lower compliance (43, 44). In many settings, preserving the supply of N95 respirators for high-risk, aerosol-generating procedures is an important consideration (45).

A systematic review of observational studies on the betacoronaviruses that cause severe acute respiratory syndrome (SARS), Middle East respiratory syndrome (MERS) and COVID-19 found that the use of face protection (including respirators and medical masks) is associated with reduced risk of infection among health workers. These studies suggested that N95 or similar respirators might be associated with greater reduction in risk than medical or 12–16-layer cotton masks. However, these studies had important

limitations (recall bias, limited information about the situations when respirators were used and limited ability to measure exposures), and very few studies included in the review evaluated the transmission risk of COVID-19 (46). Most of the studies were conducted in settings in which AGPs were performed or other high-risk settings (e.g., intensive care units or where there was exposure to infected patients and health workers were not wearing adequate PPE).

WHO continues to evaluate the evidence on the effectiveness of the use of different masks and their potential harms, risks and disadvantages, as well as their combination with hand hygiene, physical distancing of at least 1 metre and other IPC measures.

Guidance

WHO's guidance on the type of respiratory protection to be worn by health workers providing care to COVID-19 patients is based on 1) WHO recommendations on IPC for epidemic- and pandemic-prone acute respiratory infections in health care (47); 2) updated systematic reviews of randomized controlled trials on the effectiveness of medical masks compared to that of respirators for reducing the risk of clinical respiratory illness, influenza-like illness (ILI) and laboratory-confirmed influenza or viral infections. WHO guidance in this area is aligned with guidelines of other professional organizations, including the European Society of Intensive Care Medicine and the Society of Critical Care Medicine, and the Infectious Diseases Society of America (48, 49).

The WHO COVID-19 IPC GDG considered all available evidence on the modes of transmission of SARS-CoV-2 and on the effectiveness of medical mask versus respirator use to protect health workers from infection and the potential for harms such as skin conditions or breathing difficulties.

Other considerations included availability of medical masks versus respirators, cost and procurement implications and equity of access by health workers across different settings.

The majority (71%) of the GDG members confirmed their support for previous recommendations issued by WHO on 5 June 2020:

1. In the absence of aerosol generating procedures (AGPs)², WHO recommends that health workers providing care to patients with suspected or confirmed COVID-19 should wear a medical mask (in addition to other PPE that are part of droplet and contact precautions).
2. In care settings for COVID-19 patients where AGPs are performed, WHO recommends that health workers should wear a respirator (N95 or FFP2 or FFP3 standard, or equivalent) in addition to other PPE that are part of airborne and contact precautions.

In general, health workers have strong preferences about having the highest perceived protection possible to prevent COVID-19 infection and therefore may place high value on the potential benefits of respirators in settings without AGPs. WHO recommends respirators primarily for settings where AGPs are performed; however, if health workers prefer them and they are sufficiently available and cost is not an issue, they could also be used during care for COVID-19 patients in other settings. For additional guidance on PPE, including PPE

² The WHO list of AGPs includes tracheal intubation, non-invasive ventilation, tracheotomy, cardiopulmonary resuscitation, manual

ventilation before intubation, bronchoscopy, sputum induction using nebulized hypertonic saline, and dentistry and autopsy procedures.

beyond mask use by health workers, see WHO IPC guidance during health care when COVID-19 infection is suspected (3) and also WHO guidance on the rational use of PPE (45).

Exhalation valves on respirators are discouraged as they bypass the filtration function for exhaled air.

B. Guidance on the use of mask by health workers, caregivers and others based on transmission scenario

Definitions

Universal masking in health facilities is defined as the requirement for all persons (staff, patients, visitors, service providers and others) to wear a mask at all times except for when eating or drinking.

Targeted continuous medical mask use is defined as the practice of wearing a medical mask by all health workers and caregivers working in clinical areas during all routine activities throughout the entire shift.

Health workers are all people primarily engaged in actions with the primary intent of enhancing health. Examples are: nursing and midwifery professionals, doctors, cleaners, other staff who work in health facilities, social workers, and community health workers.

Evidence on universal masking in health care settings

In areas where there is community transmission or large-scale outbreaks of COVID-19, universal masking has been adopted in many hospitals to reduce the potential of transmission by health workers to patients, to other staff and anyone else entering the facility (50).

Two studies found that implementation of a universal masking policy in hospital systems was associated with decreased risk of healthcare-acquired SARS-CoV-2 infection. However, these studies had serious limitations: both were before-after studies describing a single example of a phenomenon before and after an event of interest, with no concurrent control group, and other infection control measures were not controlled for (51, 52). In addition, observed decreases in health worker infections occurred too quickly to be attributable to the universal masking policy.

Guidance

Although more research on universal masking in health settings is needed, it is the expert opinion of the majority (79%) of WHO COVID-19 IPC GDG members that universal masking is advisable in geographic settings where there is known or suspected community or cluster transmission of the SARS-CoV-2 virus.

1. In areas of known or suspected community or cluster SARS-CoV-2 transmission, universal masking should be advised in all health facilities (see Table 1).
- All health workers, including community health workers and caregivers, should wear a medical mask at all times, for any activity (care of COVID-19 or non-COVID-19 patients) and in any common area (e.g., cafeteria, staff rooms).

- Other staff, visitors, outpatients and service providers should also wear a mask (medical or non-medical) at all times
 - Inpatients are not required to wear a mask (medical or non-medical) unless physical distancing of at least 1 metre cannot be maintained (e.g., when being examined or visited at the bedside) or when outside of their care area (e.g., when being transported).
 - Masks should be changed when they become soiled, wet or damaged or if the health worker/caregiver removes the mask (e.g., for eating or drinking or caring for a patient who requires droplet/contact precautions for reasons other than COVID-19).
2. In the context of known or suspected sporadic SARS-CoV-2 virus transmission, WHO provides the following guidance:
 - Health workers, including community health workers and caregivers who work in clinical areas, should continuously wear a medical mask during routine activities throughout the entire shift, apart from when eating and drinking and changing their medical masks after caring for a patient who requires droplet/contact precautions for other reasons. In all cases, medical masks must be changed when wet, soiled, or damaged; used medical masks should be properly disposed of at the end of the shift; and new clean ones should be used for the next shift or when medical masks are changed.
 - It is particularly important to adopt the continuous use of masks in potentially high transmission risk settings including triage, family physician/general practitioner offices; outpatient departments; emergency rooms; COVID-19 designated units; haematology, oncology and transplant units; and long-term health and residential facilities.
 - Staff who do not work in clinical areas (e.g., administrative staff) do not need to wear a medical mask during routine activities if they have no exposure to patients.

Whether using masks for universal masking within health facilities or targeted continuous medical mask use throughout the entire shift, health workers should ensure the following:

- Medical mask use should be combined with other measures including frequent hand hygiene and physical distancing among health workers in shared and crowded places such as cafeterias, break rooms, and dressing rooms.
- The medical mask should be changed when wet, soiled, or damaged.
- The medical mask should not be touched to adjust it or if displaced from the face for any reason. If this happens, the mask should be safely removed and replaced, and hand hygiene performed.
- The medical mask (as well as other personal protective equipment) should be discarded and changed after caring for any patient who requires contact/droplet precautions for other pathogens, followed by hand hygiene.
- Under no circumstances should medical masks be shared between health workers or between others wearing them. Masks should be appropriately disposed of whenever removed and not reused.

- A particulate respirator at least as protective as a United States of America (US) National Institute for Occupational Safety and Health-certified N95, N99, US Food and Drug Administration surgical N95, European Union standard FFP2 or FFP3, or equivalent, should be worn in settings for COVID-19 patients where AGPs are performed (see WHO recommendations below). In these settings, this includes continuous use by health workers throughout the entire shift, when this policy is implemented.

Note: Decision makers may consider the transmission intensity in the catchment area of the health facility or community setting and the feasibility of implementing a universal masking policy compared to a policy based on assessed or presumed exposure risk. Decisions need to take into account procurement, sustainability and costs of the policy. When planning masks for all health workers, long-term availability of adequate medical masks (and when applicable, respirators) for all workers should be ensured, in particular for those providing care for patients with confirmed or suspected COVID-19. Proper use and adequate waste management should be ensured.

The potential harms and risks of mask and respirator use in the health facility setting include:

- contamination of the mask due to its manipulation by contaminated hands (53, 54);
- potential self-contamination that can occur if medical masks are not changed when wet, soiled or damaged; or by frequent touching/adjusting when worn for prolonged periods (55);
- possible development of facial skin lesions, irritant dermatitis or worsening acne, when used frequently for long hours (56-58);
- discomfort, facial temperature changes and headaches from mask wearing (44, 59, 60);
- false sense of security leading potentially to reduced adherence to well recognized preventive measures such as physical distancing and hand hygiene; and risk-taking behaviours (61-64);
- difficulty wearing a mask in hot and humid environments
- possible risk of stock depletion due to widespread use in the context of universal masking and targeted continuous mask use and consequent scarcity or unavailability for health workers caring for COVID 19 patients and during health care interactions with non-COVID-19 patients where medical masks or respirators might be required.

Alternatives to medical masks in health care settings

The WHO's disease commodity package (DCP) for COVID-19 recommends medical masks for health workers to be type II or higher (65). Type II medical masks provide a physical barrier to fluids and particulate materials and have bacterial filtration efficiency of $\geq 98\%$ compared to Type I mask, which has bacterial filtration efficiency of $\geq 95\%$ and lower fluid resistance (66) In case of stock outs of type II or higher medical masks, health workers should use a type I medical mask as an alternative. Other alternatives such as face shields or fabric masks should be carefully evaluated.

Face shields are designed to provide protection from splashes of biological fluid (particularly respiratory secretions), chemical agents and debris (67, 68) into the eyes. In the context of protection from SARS-CoV-2 transmission through respiratory droplets, face shields are used by health workers as personal protective equipment (PPE) for eye protection in combination with a medical mask or a respirator (69, 70) While a face shield may confer partial protection of the facial area against respiratory droplets, these and smaller droplets may come into contact with mucous membranes or with the eyes from the open gaps between the visor and the face (71,67).

Fabric masks are not regulated as protective masks or part of the PPE directive. They vary in quality and are not subject to mandatory testing or common standards and as such are not considered an appropriate alternative to medical masks for protection of health workers. One study that evaluated the use of cloth masks in a health care facility found that health care workers using 2 ply cotton cloth masks (a type of fabric mask) were at increased risk of influenza-like illness compared with those who wore medical masks (72).

In the context of severe medical mask shortage, face shields alone or in combination with fabric mask may be considered as a last resort (73). Ensure proper design of face shields to cover the sides of the face and below the chin.

As for other PPE items, if production of fabric masks for use in health care settings is proposed locally in situations of shortage or stock out, a local authority should assess the product according to specific minimum performance standards and required technical specifications (see Annex).

Additional considerations for community care settings

Like other health workers, community health workers should apply standard precautions for all patients at all times, with particular emphasis regarding hand and respiratory hygiene, surface and environmental cleaning and disinfection and the appropriate use of PPE. When a patient is suspected or confirmed of having COVID-19, community health workers should always apply contact and droplet precautions. These include the use of a medical mask, gown, gloves and eye protection (74).

IPC measures that are needed will depend on the local COVID-19 transmission dynamics and the type of contact required by the health care activity (see Table 1). The community health workforce should ensure that patients and workforce members apply precautionary measures such as respiratory hygiene and physical distancing of at least 1 metre (3.3 feet). They also may support set-up and maintenance of hand hygiene stations and community education (74). In the context of known or suspected community or cluster transmission, community health workers should wear a medical mask when providing essential routine services (see Table 1).

Table 1. Mask use in health care settings depending on transmission scenario, target population, setting, activity and type*

Transmission scenario	Target population (who)	Setting (where)	Activity (what)	Mask type (which one) *
Known or suspected community or cluster transmission of SARS-CoV-2	Health workers and caregivers	Health facility (including primary, secondary, tertiary care levels, outpatient care, and long-term care facilities)	For any activity in patient-care areas (COVID-19 or non-COVID-19 patients) or in any common areas (e.g., cafeteria, staff rooms)	Medical mask (or respirator if aerosol generating procedures performed)
	Other staff, patients, visitors, service suppliers		For any activity or in any common area	Medical or fabric mask
	Inpatients		When physical distance of at least 1 metre cannot be maintained	
	Health workers and caregivers	Home visit (for example, for antenatal or postnatal care, or for a chronic condition)	When in direct contact with a patient or when a distance of at least 1 metre cannot be maintained.	Medical mask
		Community	Community outreach programmes/essential routine services	
Known or suspected sporadic transmission of SARS-CoV-2 cases	Health workers and caregivers	Health facility (including primary, secondary, tertiary care levels, outpatient care, and long-term care facilities)	In patient care area- irrespective of whether patients have suspected/confirmed COVID-19	Medical mask
	Other staff, patients, visitors, service suppliers and all others		No routine activities in patient areas	Medical mask not required. Medical mask should be worn if in contact or within 1 metre of patients, or according to local risk assessment
	Health workers and caregivers	Home visit (for example, for antenatal or postnatal care, or for a chronic condition)	When in direct contact or when a distance of at least 1metre cannot be maintained.	Medical mask
		Community	Community outreach programs (e.g., bed net distribution)	
No documented SARS-CoV-2 transmission	Health workers and caregivers	Health facility (including primary, secondary, tertiary care levels, outpatient care, and long-term care facilities)	Providing any patient care	Medical mask use according to standard and transmission-based precautions
		Community	Community outreach programs	
Any transmission scenario	Health workers	Health care facility (including primary, secondary, tertiary care levels, outpatient care, and long-term care facilities), in settings where aerosol generating procedures (AGP) are performed	Performing an AGP on a suspected or confirmed COVID-19 patient or providing care in a setting where AGPs are in place for COVID-19 patients	Respirator (N95 or N99 or FFP2 or FFP3)

**This table refers only to the use of medical masks and respirators. The use of medical masks and respirators may need to be combined with other personal protective equipment and other measures as appropriate, and always with hand hygiene.*

Guidance on mask use in community settings

Evidence on the protective effect of mask use in community settings

At present there is only limited and inconsistent scientific evidence to support the effectiveness of masking of healthy people in the community to prevent infection with respiratory viruses, including SARS-CoV-2 (75). A large randomized community-based trial in which 4862 healthy participants were divided into a group wearing medical/surgical masks and a control group found no difference in infection with SARS-CoV-2 (76). A recent systematic review found nine trials (of which eight were cluster-randomized controlled trials in which clusters of people, versus individuals, were randomized) comparing medical/surgical masks versus no masks to prevent the spread of viral respiratory illness. Two trials were with healthcare workers and seven in the community. The review concluded that wearing a mask may make little or no difference to the prevention of influenza-like illness (ILI) (RR 0.99, 95%CI 0.82 to 1.18) or laboratory confirmed illness (LCI) (RR 0.91, 95%CI 0.66-1.26) (44); the certainty of the evidence was low for ILI, moderate for LCI.

By contrast, a small retrospective cohort study from Beijing found that mask use by entire families before the first family member developed COVID-19 symptoms was 79% effective in reducing transmission (OR 0.21, 0.06-0.79) (77). A case-control study from Thailand found that wearing a medical or non-medical mask all the time during contact with a COVID-19 patient was associated with a 77% lower risk of infection (aOR 0.23; 95% CI 0.09–0.60) (78). Several small observational studies with epidemiological data have reported an association between mask use by an infected person and prevention of onward transmission of SARS-CoV-2 infection in public settings. (8, 79-81).

A number of studies, some peer reviewed (82-86) but most published as pre-prints (87-104), reported a decline in the COVID-19 cases associated with face mask usage by the public, using country- or region-level data. One study reported an association between community mask wearing policy adoption and increased movement (less time at home, increased visits to commercial locations) (105). These studies differed in setting, data sources and statistical methods and have important limitations to consider (106), notably the lack of information about actual exposure risk among individuals, adherence to mask wearing and the enforcement of other preventive measures (107, 108).

Studies of influenza, influenza-like illness and human coronaviruses (not including COVID-19) provide evidence that the use of a medical mask can prevent the spread of infectious droplets from a symptomatic infected person to someone else and potential contamination of the environment by these droplets (75). There is limited evidence that wearing a medical mask may be beneficial for preventing transmission between healthy individuals sharing households with a sick person or among attendees of mass gatherings (44, 109-114).

A meta-analysis of observational studies on infections due to betacoronaviruses, with the intrinsic biases of observational data, showed that the use of either disposable medical masks or reusable 12–16-layer cotton masks was associated with protection of healthy individuals within households and among contacts of cases (46). This could be considered to be indirect evidence for the use of masks (medical or other) by healthy individuals in the wider community; however, these studies suggest that such individuals would need to be in close proximity to an infected person in a household or at a mass gathering where physical distancing cannot be achieved to become infected with the virus. Results from cluster randomized controlled trials on the use of masks among young adults living in university residences in the United States of America indicate that face masks may reduce the rate of influenza-like illness but showed no impact on risk of laboratory-confirmed influenza (115, 116).

Guidance

The WHO COVID-19 IPC GDG considered all available evidence on the use of masks by the general public including effectiveness, level of certainty and other potential benefits and harms, with respect to transmission scenarios, indoor versus outdoor settings, physical distancing and ventilation. Despite the limited evidence of protective efficacy of mask wearing in community settings, in addition to all other recommended preventive measures, the GDG advised mask wearing in the following settings:

1. In areas with known or suspected community or cluster transmission of SARS-CoV-2, WHO advises mask use by the public in the following situations (see Table 2):

Indoor settings:

- in public indoor settings where ventilation is known to be poor regardless of physical distancing: limited or no opening of windows and doors for natural ventilation; ventilation system is not properly functioning or maintained; or cannot be assessed;
- in public indoor settings that have adequate³ ventilation if physical distancing of at least 1 metre cannot be maintained;
- in household indoor settings: when there is a visitor who is not a household member and ventilation is known to be poor, with limited opening of windows and doors for natural ventilation, or the ventilation system cannot be assessed or is not properly functioning, regardless of whether physical distancing of at least 1 metre can be maintained;
- in household indoor settings that have adequate ventilation if physical distancing of at least 1 metre cannot be maintained.

³ For adequate ventilation refer to regional or national institutions or heating, refrigerating and air-conditioning societies enacting ventilation requirements. If not available or applicable, a recommended ventilation rate of 10 l/s/person should be met (except healthcare facilities which have specific requirements). For more information consult “Coronavirus (COVID-19) response

resources from ASHRAE and others”
<https://www.ashrae.org/technical-resources/resources>

Table 2. Mask use in community settings depending on transmission scenario, setting, target population, purpose and type*

Transmission scenario	Situations/settings (where)	Target Population (who)	Purpose of mask use (why)	Mask type (which one)
Known or suspected community or cluster transmission of SARS-CoV-2	Indoor settings, where ventilation is known to be poor or cannot be assessed or the ventilation system is not properly maintained, regardless of whether physical distancing of at least 1 meter can be maintained	General population in public* settings such as shops, shared workplaces, schools, churches, restaurants, gyms, etc. or in enclosed settings such as public transportation.	Potential benefit for source control	Fabric mask
	Indoor settings that have adequate ⁴ ventilation if physical distancing of at least 1 metre cannot be maintained	For households, in indoor settings, when there is a visitor who is not a member of the household		
	Outdoor settings where physical distancing cannot be maintained	General population in settings such as crowded open-air markets, lining up outside a building, during demonstrations, etc.		
	Settings where physical distancing cannot be maintained, and the individual is at increased risk of infection and/or negative outcomes	Individuals/people with higher risk of severe complications from COVID-19: • People aged ≥ 60 years • People with underlying comorbidities, such as cardiovascular disease or diabetes mellitus, chronic lung disease, cancer, cerebrovascular disease, immunosuppression, obesity, asthma	Protection	Medical mask
Known or suspected sporadic transmission, or no documented SARS-CoV-2 transmission	Risk-based approach	General population	Potential benefit for source control and/or protection	Depends on purpose (see details in the guidance content)
Any transmission scenario	Any setting in the community	Anyone suspected or confirmed of having COVID-19, regardless of whether they have symptoms or not, or anyone awaiting viral test results, when in the presence of others	Source control	Medical mask

*Public indoor setting includes any indoor setting outside of the household

⁴ For adequate ventilation refer to regional or national institutions or heating, refrigerating and air-conditioning societies enacting ventilation requirements. If not available or applicable, a recommended ventilation rate of 10l/s/person should be met (except healthcare facilities which have specific requirements).). For more information consult “Coronavirus (COVID-19) response resources from ASHRAE and others” <https://www.ashrae.org/technical-resources/resources>

In outdoor settings:

- where physical distancing of at least 1 metre cannot be maintained;
- individuals/people with higher risk of severe complications from COVID-19 (individuals ≥ 60 years old and those with underlying conditions such as cardiovascular disease or diabetes mellitus, chronic lung disease, cancer, cerebrovascular disease or immunosuppression) should wear medical masks in any setting where physical distance cannot be maintained.

2. In areas with known or suspected sporadic transmission or no documented transmission, as in all transmission scenarios, WHO continues to advise that decision makers should apply a risk-based approach focusing on the following criteria when considering the use of masks for the public:

- **Purpose of mask use.** Is the intention source control (preventing an infected person from transmitting the virus to others) or protection (preventing a healthy wearer from the infection)?
- **Risk of exposure to SARS-CoV-2.** Based on the epidemiology and intensity of transmission in the population, is there transmission and limited or no capacity to implement other containment measures such as contact tracing, ability to carry out testing and isolate and care for suspected and confirmed cases? Is there risk to individuals working in close contact with the public (e.g., social workers, personal support workers, teachers, cashiers)?
- **Vulnerability of the mask wearer/population.** Is the mask wearer at risk of severe complications from COVID-19? Medical masks should be used by older people (≥ 60 years old), immunocompromised patients and people with comorbidities, such as cardiovascular disease or diabetes mellitus, chronic lung disease, cancer and cerebrovascular disease (117).
- **Setting in which the population lives.** Is there high population density (such as in refugee camps, camp-like settings, and among people living in cramped conditions) and settings where individuals are unable to keep a physical distance of at least 1 metre (for example, on public transportation)?
- **Feasibility.** Are masks available at an affordable cost? Do people have access to clean water to wash fabric masks, and can the targeted population tolerate possible adverse effects of wearing a mask?
- **Type of mask.** Does the use of medical masks in the community divert this critical resource from the health workers and others who need them the most? In settings where medical masks are in short supply, **stocks should be prioritized for health workers and at-risk individuals.**

The decision of governments and local jurisdictions whether to recommend or make mandatory the use of masks should be based on the above assessment as well as the local context, culture, availability of masks and resources required.

3. In any transmission scenario:

- Persons with any symptoms suggestive of COVID-19 should wear a medical mask and (5) additionally:
 - self-isolate and seek medical advice as soon as they start to feel unwell with potential symptoms of COVID-19, even if symptoms are mild);

- follow instructions on how to put on, take off, and dispose of medical masks and perform hand hygiene (118);
- follow all additional measures, in particular respiratory hygiene, frequent hand hygiene and maintaining physical distance of at least 1 metre from other persons (46). If a medical mask is not available for individuals with suspected or confirmed COVID-19, a fabric mask meeting the specifications in the Annex of this document should be worn by patients as a source control measure, pending access to a medical mask. The use of a non-medical mask can minimize the projection of respiratory droplets from the user (119, 120).
- Asymptomatic persons who test positive for SARS-CoV-2, should wear a medical mask when with others for a period of 10 days after testing positive.

Potential benefits/harms

The potential advantages of mask use by healthy people in the general public include:

- reduced spread of respiratory droplets containing infectious viral particles, including from infected persons before they develop symptoms (121);
- reduced potential for stigmatization and greater of acceptance of mask wearing, whether to prevent infecting others or by people caring for COVID-19 patients in non-clinical settings (122);
- making people feel they can play a role in contributing to stopping spread of the virus;
- encouraging concurrent transmission prevention behaviours such as hand hygiene and not touching the eyes, nose and mouth (123-125);
- preventing transmission of other respiratory illnesses like tuberculosis and influenza and reducing the burden of those diseases during the pandemic (126).

The potential disadvantages of mask use by healthy people in the general public include:

- headache and/or breathing difficulties, depending on type of mask used (55);
- development of facial skin lesions, irritant dermatitis or worsening acne, when used frequently for long hours (58, 59, 127);
- difficulty with communicating clearly, especially for persons who are deaf or have poor hearing or use lip reading (128, 129);
- discomfort (44, 55, 59)
- a false sense of security leading to potentially lower adherence to other critical preventive measures such as physical distancing and hand hygiene (105);
- poor compliance with mask wearing, in particular by young children (111, 130-132);
- waste management issues; improper mask disposal leading to increased litter in public places and environmental hazards (133);
- disadvantages for or difficulty wearing masks, especially for children, developmentally challenged persons, those with mental illness, persons with cognitive impairment, those with asthma or chronic respiratory or breathing problems, those who have had facial trauma or recent oral maxillofacial surgery and those living in hot and humid environments (55, 130).

Considerations for implementation

When implementing mask policies for the public, decision-makers should:

- clearly communicate the purpose of wearing a mask, including when, where, how and what type of mask should be worn; explain what wearing a mask may achieve and what it will not achieve; and communicate clearly that this is one part of a package of measures along with hand hygiene, physical distancing, respiratory etiquette, adequate ventilation in indoor settings and other measures that are all necessary and all reinforce each other;
- inform/train people on when and how to use masks appropriately and safely (see mask management and maintenance sections);
- consider the feasibility of use, supply/access issues (cleaning, storage), waste management, sustainability, social and psychological acceptance (of both wearing and not wearing different types of masks in different contexts);
- continue gathering scientific data and evidence on the effectiveness of mask use (including different types of masks) in non-health care settings;
- evaluate the impact (positive, neutral or negative) of using masks in the general population (including behavioural and social sciences) through good quality research.

Mask use during physical activity

Evidence

There are limited studies on the benefits and harms of wearing medical masks, respirators and non-medical masks while exercising. Several studies have demonstrated statistically significant deleterious effects on various cardiopulmonary physiologic parameters during mild to moderate exercise in healthy subjects and in those with underlying respiratory diseases (134-140). The most significant impacts have been consistently associated with the use of respirators and in persons with underlying obstructive airway pulmonary diseases such as asthma and chronic obstructive pulmonary disease (COPD), especially when the condition is moderate to severe (136). Facial microclimate changes with increased temperature, humidity and perceptions of dyspnoea were also reported in some studies on the use of masks during exercise (134, 141). A recent review found negligible evidence of negative effects of mask use during exercise but noted concern for individuals with severe cardiopulmonary disease (142).

Guidance

WHO advises that people should not wear masks during vigorous intensity physical activity (143) because masks may reduce the ability to breathe comfortably. The most important preventive measure is to maintain physical distancing of at least 1 meter and ensure good ventilation when exercising.

If the activity takes place indoors, adequate ventilation should be ensured at all times through natural ventilation or a properly functioning or maintained ventilation system (144). Particular attention should be paid to cleaning and disinfection of the environment, especially high-touch surfaces. If all the above measures cannot be ensured, consider temporary closure of public indoor exercise facilities (e.g., gyms).

Face shields for the general public

At present, face shields are considered to provide a level of eye protection only and should not be considered as an equivalent to masks with respect to respiratory droplet protection and/or source control. Current laboratory testing standards only assess face shields for their ability to provide eye protection from chemical splashes (145).

In the context of non-availability or difficulties wearing a non-medical mask (in persons with cognitive, respiratory or hearing impairments, for example), face shields may be considered as an alternative, noting that they are inferior to masks with respect to droplet transmission and prevention. If face shields are to be used, ensure proper design to cover the sides of the face and below the chin.

Medical masks for the care of COVID-19 patients at home

WHO provides guidance on how to care for patients with confirmed and suspected COVID-19 at home when care in a health facility or other residential setting is not possible (5).

- Persons with suspected COVID-19 or mild COVID-19 symptoms should wear a medical mask as much as possible, especially when there is no alternative to being in the same room with other people. The mask should be changed at least once daily. Persons who cannot tolerate a medical mask should rigorously apply respiratory hygiene (i.e., cover mouth and nose with a disposable paper tissue when coughing or sneezing and dispose of it immediately after use or use a bent elbow procedure and then perform hand hygiene).
- Caregivers of or those sharing living space with people with suspected COVID-19 or with mild COVID-19 symptoms should wear a medical mask when in the same room as the affected person.

References

1. World Health Organization. The World Health Report 2006 - working together for health. Geneva: World Health Organization; 2006. (<https://apps.who.int/iris/handle/10665/43432> accessed 21 November 2020)
2. World Health Organization. Advice on the use of masks for children in the context of COVID-19. Annex to the Advice on the use of masks in the context of COVID-19. Geneva, 2020. (<https://apps.who.int/iris/handle/10665/333919> accessed 21 November 2020).
3. World Health Organization. Infection prevention and control during health care when COVID-19 is suspected: interim guidance. Geneva: World Health Organization; 2020. (<https://apps.who.int/iris/handle/10665/332879> accessed 21 November 2020).
4. World Health Organization. Infection prevention and control for long-term care facilities in the context of COVID-19: interim guidance. Geneva: World Health Organization; 2020. (<https://apps.who.int/iris/handle/10665/331508> accessed 21 November 2020).

5. World Health Organization. Home care for patients with suspected or confirmed COVID-19 and management of their contacts: interim guidance. Geneva: World Health Organization; 2020 (<https://apps.who.int/iris/handle/10665/333782>. accessed 21 November 2020).
6. World Health Organization. Infection prevention and control of epidemic-and pandemic prone acute respiratory infections in health care. (https://apps.who.int/iris/bitstream/handle/10665/112656/9789241507134_eng.pdf?sequence=1 Accessed 21 November 2020).
7. Machida M, Nakamura I, Saito R, Nakaya T, Hanibuchi T, Takamiya T, et al. Incorrect Use of Face Masks during the Current COVID-19 Pandemic among the General Public in Japan. *Int J Environ Res Public Health*. 2020;17(18).
8. Liu J, Liao X, Qian S, Yuan J, Wang F, Liu Y, et al. Community Transmission of Severe Acute Respiratory Syndrome Coronavirus 2, Shenzhen, China, 2020. *Emerg Infect Dis*. 2020;26(6):1320-3.
9. Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*. 2020;395(10223):514-23.
10. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506.
11. Burke RM, Midgley CM, Dratch A, Fenstersheib M, Haupt T, Holshue M, et al. Active Monitoring of Persons Exposed to Patients with Confirmed COVID-19 - United States, January-February 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(9):245-6.
12. World Health Organization. Transmission of SARS-CoV-2: implications for infection prevention precautions. (<https://apps.who.int/iris/handle/10665/333114> accessed 21 November 2020).
13. World Health Organization. Coronavirus disease 2019 (COVID-19) Situation Report – 73. Geneva: World Health Organization; 2020 (https://www.who.int/docs/default-source/coronavirus/situation-reports/20200402-sitrep-73-covid-19.pdf?sfvrsn=5ae25bc7_6, accessed 21 November 2020).
14. Cheng VCC, Wong SC, Chen JHK, Yip CCY, Chuang VWM, Tsang OTY, et al. Escalating infection control response to the rapidly evolving epidemiology of the coronavirus disease 2019 (COVID-19) due to SARS-CoV-2 in Hong Kong. *Infect Control Hosp Epidemiol*. 2020;41(5):493-8.
15. Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, Surface Environmental, and Personal Protective Equipment Contamination by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) From a Symptomatic Patient. *JAMA*. 2020. 323(16):1610-1612. doi: 10.1001/jama.2020.3227.
16. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med*. 2020;382(16):1564-7.
17. Meyerowitz EA, Richterman A, Gandhi RT, Sax PE. Transmission of SARS-CoV-2: A Review of Viral, Host, and Environmental Factors. *Ann Intern Med*. 2020;M20-5008. doi:10.7326/M20-5008
18. Wei J, Li Y. Airborne spread of infectious agents in the indoor environment. *Am J Infect Control*. 2016;44(9 Suppl):S102-8.
19. McCarthy J, McCarthy M. Long range versus short range aerial transmission of SARS-CoV-2. 2020 (<https://arxiv.org/pdf/2008.03558.pdf> (Accessed 24 November 2020)).
20. Lednicky JA, Lauzardo M, Fan ZH, et al. Viable SARS-CoV-2 in the air of a hospital room with COVID-19 patients. *medRxiv*. doi:10.1101/2020.08.03.20167395
21. Ring N, Ritchie K, Mandava L, Jepson R. A guide to synthesising qualitative research for researchers undertaking health technology assessments and systematic reviews. 2011. NHS Quality Improvement Scotland (NHS QIS).
22. Wolfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Muller MA, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature*. 2020;581(7809):465-9.
23. van Kampen J, van de Vijver D, Fraaij P, Haagmans B, Lamers M, Okba Nea. Shedding of infectious virus in hospitalized patients with coronavirus disease-2019 (COVID19): duration and key determinants. *MedRxiv*. 2020 doi:10.1101/2020.06.08.20125310.
24. Centers for Disease Control and Prevention. Symptom-Based Strategy to Discontinue Isolation for Persons with COVID-19. Atlanta: Centers for Disease Control and Prevention; (<https://www.cdc.gov/coronavirus/2019-ncov/community/strategy-discontinue-isolation.html>, accessed 21 November 2020).
25. Yu P, Zhu J, Zhang Z, Han Y. A Familial Cluster of Infection Associated With the 2019 Novel Coronavirus Indicating Possible Person-to-Person Transmission During the Incubation Period. *J Infect Dis*. 2020;221(11):1757-61. E
26. Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern Med*. 2020;172(9):577-82.
27. Kimball A, Hatfield KM, Arons M, James A, Taylor J, Spicer K, et al. Asymptomatic and Presymptomatic SARS-CoV-2 Infections in Residents of a Long-Term Care Skilled Nursing Facility - King County, Washington, March 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(13):377-81.
28. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*. 2020;26(5):672-5.
29. Buitrago-Garcia D, Egli-Gany D, Counotte MJ, Hossmann S, Imeri H, Ipekci AM, et al. Occurrence and transmission potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis. *PLoS Med*. 2020;17(9):e1003346.
30. Byambasuren O, Cardona, M., Bell, K., Clark, J., McLaws, M.-L., Glasziou, P. Estimating the extent of true asymptomatic COVID-19 and its potential for community transmission: systematic review and meta-analysis. *JAMMI* 2020 doi: 10.3138/jammi-2020-0030

31. Arons MM, Hatfield KM, Reddy SC, Kimball A, James A, Jacobs JR, et al. Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled Nursing Facility. *N Engl J Med*. 2020;382(22):2081-90.
32. Hu Z, Song C, Xu C, Jin G, Chen Y, Xu X, et al. Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci*. 2020;63(5):706-11.
33. Huang R, Xia J, Chen Y, Shan C, Wu C. A family cluster of SARS-CoV-2 infection involving 11 patients in Nanjing, China. *Lancet Infect Dis*. 2020;20(5):534-5.
34. Pan X, Chen D, Xia Y, Wu X, Li T, Ou X, et al. Asymptomatic cases in a family cluster with SARS-CoV-2 infection. *Lancet Infect Dis*. 2020;20(4):410-1.
35. Wang Y, Tong J, Qin Y, Xie T, Li J, Li J, et al. Characterization of an asymptomatic cohort of SARS-CoV-2 infected individuals outside of Wuhan, China. *Clin Infect Dis*. 2020; 71(16):2132-2138. doi: 10.1093/cid/ciaa629.
36. Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic Transmission of SARS-CoV-2 - Singapore, January 23-March 16, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(14):411-5.
37. Bae SH, Shin H, Koo HY, Lee SW, Yang JM, Yon DK. Asymptomatic Transmission of SARS-CoV-2 on Evacuation Flight. *Emerg Infect Dis*. 2020;26(11).
38. Qiu X, Nergiz I, Maraolo A, Bogoch, Low N, Cevik M. Defining the role of asymptomatic SARS-CoV-2 transmission: a living systematic review. *MedRxiv*. 2020 doi: 10.1101/2020.09.01.20135194.
39. European Standards. UNE EN 14683:2019+AC:2019. Medical Face Masks -Requirements and Test Methods. 2019; (<https://www.en-standard.eu/une-en-14683-2019-ac-2019-medical-face-masks-requirements-and-test-methods/> accessed 21 November 2020)
40. ASTM International. F23 Committee. Specification for Performance of Materials Used in Medical Face Masks. (<https://www.astm.org/Standards/F2100.htm> accessed 21 November 2020)
41. National Institute for Occupational Safety and Health (NIOSH). NIOSH Guide to the Selection and Use of Particulate Respirators. Department of Health and Human Services (DHHS) NIOSH publication number 96-101, 1996. (<https://www.cdc.gov/niosh/docs/96-101/default.html> accessed 21 November 2020)
42. CEN, E., 2001. 149: 2001 norm: Respiratory protective devices-Filtering half masks to protect against particles-Requirements, testing, marking. European Committee for Standardization. (<https://shop.bsigroup.com/ProductDetail?pid=0000000000030178264> accessed 21 November 2020).
43. Long Y, Hu T, Liu L, Chen R, Guo Q, Yang L, et al. Effectiveness of N95 respirators versus surgical masks against influenza: A systematic review and meta-analysis. *J Evid Based Med*. 2020;13(2):93-101.
44. Jefferson T DMC, Dooley L, Ferroni E, Al-Ansary LA, Bawazeer GA, et al. Physical interventions to interrupt or reduce the spread of respiratory viruses. *Cochrane Database of Systematic Reviews* 2020;(11):CD006207. doi:10.1002/14651858.CD006207.pub5. 2020.
45. World Health Organization. Rational use of personal protective equipment for coronavirus disease 2019 (COVID-19). Geneva: World Health Organization; 2020 (<https://apps.who.int/iris/rest/bitstreams/1274340/retrieve> accessed 21 November 2020).
46. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schunemann HJ, et al. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet*. 2020;395(10242):1973-87.
47. Chan AJ, Islam MK, Rosewall T, Jaffray DA, Easty AC, Cafazzo JA. The use of human factors methods to identify and mitigate safety issues in radiation therapy. *Radiotherapy and Oncology*. 2010;97(3):596-600.
48. Brilli RJ, Spevetz A, Branson RD, Campbell GM, Cohen H, Dasta JF, et al. Critical care delivery in the intensive care unit: defining clinical roles and the best practice model. *Crit Care Med*. 2001;29(10):2007-19.
49. Roland D, McCaffery K, Davies F. Scoring systems in paediatric emergency care: Panacea or paper exercise? *Journal of paediatrics and child health*. 2016;52(2):181-6.
50. Klompas M, Morris CA, Sinclair J, Pearson M, Shenoy ES. Universal Masking in Hospitals in the Covid-19 Era. *N Engl J Med*. 2020;382(21):e63
51. Seidelman J, Lewis S, Advani S, Akinboyo I, Epling C, Case M, et al. Universal Masking is an Effective Strategy to Flatten the SARS-2-CoV Healthcare Worker Epidemiologic Curve. *Infect Control Hosp Epidemiol*. 2020:1-5.
52. Wang X, Ferro EG, Zhou G, Hashimoto D, Bhatt DL. Association Between Universal Masking in a Health Care System and SARS-CoV-2 Positivity Among Health Care Workers. *JAMA*. 2020; 324(7):703-704.
53. Zamora JE, Murdoch J, Simchison B, Day AG. Contamination: a comparison of 2 personal protective systems. *CMAJ*. 2006;175(3):249-54.
54. Kwon JH, Burnham CD, Reske KA, Liang SY, Hink T, Wallace MA, et al. Assessment of Healthcare Worker Protocol Deviations and Self-Contamination During Personal Protective Equipment Donning and Doffing. *Infect Control Hosp Epidemiol*. 2017;38(9):1077-83.
55. Bakhit M, Krzyzaniak N, Scott A, Clark J, Glasziou P, Del Mar C. Downsides of face masks and possible mitigation strategies: a systematic review and meta-analysis. *MedRxiv*. 2020 doi: 10.1101/2020.06.16.20133207.
56. Foo CC, Goon AT, Leow YH, Goh CL. Adverse skin reactions to personal protective equipment against severe acute respiratory syndrome--a descriptive study in Singapore. *Contact Dermatitis*. 2006;55(5):291-4.
57. Radonovich LJ, Jr., Simberloff MS, Bessesen MT, Brown AC, Cummings DAT, Gaydos CA, et al. N95 Respirators vs Medical Masks for Preventing Influenza Among Health Care Personnel: A Randomized Clinical Trial. *JAMA*. 2019;322(9):824-33.
58. Al Badri F. Surgical mask contact dermatitis and epidemiology of contact dermatitis in healthcare workers. *Current Allergy & Clinical Immunology*, 2017; 30,3: 183 - 188.

59. Matusiak L, Szepietowska M, Krajewski P, Bialynicki-Birula R, Szepietowski JC. Inconveniences due to the use of face masks during the COVID-19 pandemic: a survey study of 876 young people. *Dermatol Ther*. 2020. doi: 10.1111/dth.13567
60. MacIntyre CR, Wang Q, Cauchemez S, Seale H, Dwyer DE, Yang P, et al. A cluster randomized clinical trial comparing fit-tested and non-fit-tested N95 respirators to medical masks to prevent respiratory virus infection in health care workers. *Influenza Other Respir Viruses*. 2011;5(3):170-9.
61. Morrongiello BA, Major K. Influence of safety gear on parental perceptions of injury risk and tolerance or children's risk taking. *Inj Prev*. 2002;8(1):27-31.
62. Morrongiello BA, Walpole B, Lasenby J. Understanding children's injury-risk behavior: wearing safety gear can lead to increased risk taking. *Accid Anal Prev*. 2007;39(3):618-23.
63. Lasenby-Lessard J, Morrongiello BA. Understanding risk compensation in children: Experience with the activity and level of sensation seeking play a role. *Accid Anal Prev*. 2011;43(4):1341-7.
64. DiLillo D, Tremblay G. Maternal and child reports of behavioral compensation in response to safety equipment usage. *J Pediatr Psychol*. 2001;26(3):175-84.
65. Thomas EJ, Sexton JB, Helmreich RL. Translating teamwork behaviours from aviation to healthcare: development of behavioural markers for neonatal resuscitation. *Qual Saf Health Care*. 2004 Oct; 13(Suppl 1): i57-i64.
66. Pri-Med Medicinal Products. Mask Protection Standards & Medical Face Mask Information For Use. (<https://www.primed.ca/clinical-resources/astm-mask-protection-standards/#:~:text=Are%20there%20different%20levels%20of%20protection%20with%20ASTM-rated,%20%20160%20%201%20more%20rows%20> accessed 21 November 2020).
67. Hirschmann MT, Hart A, Henckel J, Sadoghi P, Seil R, Mouton C. COVID-19 coronavirus: recommended personal protective equipment for the orthopaedic and trauma surgeon. *Knee Surg Sports Traumatol Arthrosc*. 2020;28(6):1690-8.
68. Anon JB, Denne C, Rees D. Patient-Worn Enhanced Protection Face Shield for Flexible Endoscopy. *Otolaryngol Head Neck Surg*. 2020;163(2):280-3.
69. McBride ME, Waldrop WB, Fehr JJ, Boulet JR, Murray DJ. Simulation in pediatrics: the reliability and validity of a multiscenario assessment. *Pediatrics*. 2011; 128: 335-343.
70. Kähler CJ, Hain R. Fundamental protective mechanisms of face masks against droplet infections. *J Aerosol Sci*. 2020; 148: 105617.
71. Lindsley WG, Noti JD, Blachere FM, Szalajda JV, Beezhold DH. Efficacy of face shields against cough aerosol droplets from a cough simulator. *J Occup Environ Hyg*. 2014;11(8):509-18.
72. MacIntyre CR, Seale H, Dung TC, Hien NT, Nga PT, Chughtai AA, et al. A cluster randomised trial of cloth masks compared with medical masks in healthcare workers. *BMJ Open*. 2015;5(4):e006577.
73. Centers for Disease Control and Prevention. If You Are Immunocompromised, Protect Yourself From COVID-19. (<https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/immunocompromised.html> accessed 21 November 2020).
74. Nielsen J, Landauer TK, editors. A mathematical model of the finding of usability problems. *Proceedings of the INTERACT'93 and CHI'93 conference on Human factors in computing systems*. ACM. 1993; 206-213.
75. Chou R, Dana T, Jungbauer R, Weeks C, McDonagh MS. Masks for Prevention of Respiratory Virus Infections, Including SARS-CoV-2, in Health Care and Community Settings: A Living Rapid Review. *Ann Intern Med*. 2020;173(7):542-555. doi:10.7326/M20-3213
76. Bundgaard H, J. B, Raaschou-Pedersen D, von Buchwald C, Todsén T, Norsk J. Effectiveness of Adding a Mask Recommendation to Other Public Health Measures to Prevent SARS-CoV-2 Infection in Danish Mask Wearers. *Ann Intern Med*. 2020. doi: 10.7326/M20-6817.
77. Wang Y, Tian H, Zhang L, Zhang M, Guo D, Wu W, et al. Reduction of secondary transmission of SARS-CoV-2 in households by face mask use, disinfection and social distancing: a cohort study in Beijing, China. *BMJ Glob Health*. 2020; 5(5): e002794.
78. Doung-ngern P, Suphanchaimat R, Panjangampatthana A, Janekrongtham C, Ruampoom D, Daochaeng N. Associations between mask-wearing, handwashing, and social distancing practices and risk of COVID-19 infection in public: a case-control study in Thailand. *Emerg Infect Dis*. 2020;26(11):2607-2616.
79. Chen J, He H, Cheng W. Potential transmission of SARS-CoV-2 on a flight from Singapore to Hangzhou, China: An epidemiological investigation. *Travel Med Infect Dis*. 2020; 36: 101816.
80. Hendrix MJ, Walde C, Findley K, Trotman R. Absence of Apparent Transmission of SARS-CoV-2 from Two Stylists After Exposure at a Hair Salon with a Universal Face Covering Policy - Springfield, Missouri, May 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(28):930-2.
81. Schwartz KL, Murti M, Finkelstein M, Leis JA, Fitzgerald-Husek A, Bourns L, et al. Lack of COVID-19 transmission on an international flight. *CMAJ*. 2020;192(15):E410.
82. Chiang CH, Chiang CH, Chiang CH, Chen YC. The Practice of Wearing Surgical Masks during the COVID-19 Pandemic. *Emerg Infect Dis*. 2020;26(8):1962.
83. Cheng VC, Wong SC, Chuang VW, So SY, Chen JH, Sridhar S, et al. The role of community-wide wearing of face mask for control of coronavirus disease 2019 (COVID-19) epidemic due to SARS-CoV-2. *J Infect*. 2020;81(1):107-14.
84. Bo Y, Guo C, Lin C, et al. Effectiveness of non-pharmaceutical interventions on COVID-19 transmission in 190 countries from 23 January to 13 April 2020. *Int J Infect Dis*. 2020; 102: 247-253.
85. Lyu W, Wehby GL. Community Use Of Face Masks And COVID-19: Evidence From A Natural Experiment Of State Mandates In The US. *Health Aff (Millwood)*. 2020;39(8):1419-25.

86. Gallaway MS, Rigler J, Robinson S, Herrick K, Livar E, Komatsu KK, et al. Trends in COVID-19 Incidence After Implementation of Mitigation Measures - Arizona, January 22-August 7, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(40):1460-3.
87. Rader B, White LF, Burns MR, Chen J, Brilliant J, Cohen J, et al. Mask Wearing and Control of SARS-CoV-2 Transmission in the United States. *MedRxiv.* 2020. doi: 10.1101/2020.08.23.20078964.
88. Matzinger P, Skinner J. Strong impact of closing schools, closing bars and wearing masks during the Covid-19 pandemic: results from a simple and revealing analysis. *MedRxiv.* 2020. doi: 10.1101/2020.09.26.20202457.
89. Kenyon C. Widespread use of face masks in public may slow the spread of SARS CoV-2: 1 an ecological study. *MedRxiv.* 2020. doi: 10.1101/2020.03.31.20048652.
90. Leffler CT, Ing E, Lykins JD, Hogan MC, McKeown CA, Grzybowski A. Association of Country-wide Coronavirus Mortality with Demographics, Testing, Lockdowns, and Public Wearing of Masks. *Am J Trop Med Hyg.* 2020. doi: 10.4269/ajtmh.20-1015.
91. Lan F-Y, Christophi C, Buley J, Iliaki E, Bruno-Murtha L, Sayah A, et al. Effects of universal masking on Massachusetts healthcare workers' COVID-19 incidence. *MedRxiv.* 2020. doi: 10.1101/2020.08.09.20171173.
92. Aravindakshan A, Boehnke J, Gholami E, Nayak A. Mask-Wearing During the COVID-19 Pandemic. *MedRxiv.* 2020. doi: 10.1101/2020.09.11.20192971.
93. Pletz M, Steiner A, Kesselmeier M, Loeffler B, Trommer S, Weis S, et al. Impact of universal masking in health care and community on SARS-CoV-2 spread. *MedRxiv.* 2020. doi: 10.1101/2020.09.02.20187021.
94. Fortaleza C, et al. Impact of nonpharmaceutical governmental strategies for prevention and control of COVID-19 in São Paulo State, Brazil. *MedRxiv.* 2020. doi: 10.1101/2020.08.23.20180273.
95. Karaivanov A, Lu SE, Shigeoka H, Chen C, Pamplona S. Face Masks, Public Policies and Slowing the Spread of COVID-19: Evidence from Canada. *MedRxiv.* 2020. doi: 10.1101/2020.09.24.20201178.
96. Miyazawa D, Kaneko G. Face mask wearing rate predicts country's COVID-19 death rates: with supplementary state-by-state data in the United States. *MedRxiv.* 2020. doi: 10.1101/2020.06.22.20137745.
97. Mitze T, Kosfeld R, Rode J, Walde K. Face Masks Considerably Reduce Covid-19 Cases in Germany. *MedRxiv.* 2020. doi: 10.1101/2020.06.21.20128181.
98. Maloney M, Rhodes N, Yarnold P. Mask mandates can limit COVID spread: Quantitative assessment of month-over-month effectiveness of governmental policies in reducing the number of new COVID-19 cases in 37 US States and the District of Columbia. *MedRxiv.* 2020. doi: 10.1101/2020.10.06.20208033.
99. Sruthi C, Biswal M, Saraswat B, Joshi H, Prakash M. How Policies on Restaurants, Bars, Nightclubs, Masks, Schools, and Travel Influenced Swiss COVID-19 Reproduction Ratios. *MedRxiv.* 2020. doi: 10.1101/2020.10.11.20210641.
100. Lan F, Christophi C, Buley J, Iliaki E, Bruno-Murtha L, Sayah A, et al. Effects of universal masking on Massachusetts healthcare workers' COVID-19 incidence. *MedRxiv.* 2020. doi: 10.1101/2020.08.09.20171173.
101. Shacham e, Scroggins S, Ellis M, Garza A. Association of County-Wide Mask Ordinances with Reductions in Daily CoVID-19 Incident Case Growth in a Midwestern Region Over 12 Weeks. *MedRxiv.* 2020. doi: 10.1101/2020.10.28.20221705.
102. Chernozhukov V, Kasahara H, Schrimpf P. Causal Impact of Masks, Policies, Behavior on Early Covid-19 Pandemic in the U.S. *J Econom.* 2020. doi: 10.1016/j.jeconom.2020.09.003.
103. Research GS. Face Masks and GDP. 2020. (<https://www.goldmansachs.com/insights/pages/face-masks-and-gdp.html> accessed 21 November 2020).
104. Scott N, Saul A, Spelman T, Stooze M, Pedrana A, Saeri A. The introduction of a mandatory mask policy was associated with significantly reduced COVID-19 cases in a major metropolitan city. 2020. (Available at SSRN:<http://dx.doi.org/10.2139/ssrn.3714648> accessed 29 November 2020).
105. Yan Y, Bayham J, Fenichel E, Richter A. Do Face Masks Create a False Sense of Security? A COVID-19 Dilemma. *MedRxiv.* 2020. doi: 10.1101/2020.05.23.20111302.
106. Piantadosi S, Byar DP, Green SB. The ecological fallacy. *Am J Epidemiol.* 1988;127(5):893-904.
107. Clifford GD, Long WJ, Moody GB, Szolovits P. Robust parameter extraction for decision support using multimodal intensive care data. *Philos Trans A Math Phys Eng Sci.* 2009 Jan 28; 367(1887): 411-429.
108. Dufault B, Klar N. The quality of modern cross-sectional ecologic studies: a bibliometric review. *Am J Epidemiol.* 2011;174(10):1101-7.
109. Barasheed O, Alfelali M, Mushta S, Bokhary H, Alshehri J, Attar AA, et al. Uptake and effectiveness of facemask against respiratory infections at mass gatherings: a systematic review. *Int J Infect Dis.* 2016;47:105-11.
110. Barasheed O, Almasri N, Badahdah AM, Heron L, Taylor J, McPhee K, et al. Pilot Randomised Controlled Trial to Test Effectiveness of Facemasks in Preventing Influenza-like Illness Transmission among Australian Hajj Pilgrims in 2011. *Infect Disord Drug Targets.* 2014;14(2):110-6.
111. Cowling BJ, Chan KH, Fang VJ, Cheng CK, Fung RO, Wai W, et al. Facemasks and hand hygiene to prevent influenza transmission in households: a cluster randomized trial. *Ann Intern Med.* 2009;151(7):437-46.
112. Lau JT, Tsui H, Lau M, Yang X. SARS transmission, risk factors, and prevention in Hong Kong. *Emerg Infect Dis.* 2004;10(4):587-92.
113. Suess T, Remschmidt C, Schink SB, Schweiger B, Nitsche A, Schroeder K, et al. The role of facemasks and hand hygiene in the prevention of influenza transmission in households: results from a cluster randomised trial; Berlin, Germany, 2009-2011. *BMC Infect Dis.* 2012;12:26.
114. Wu J, Xu F, Zhou W, Feikin DR, Lin CY, He X, et al. Risk factors for SARS among persons without known contact with SARS patients, Beijing, China. *Emerg Infect Dis.* 2004;10(2):210-6.

115. Aiello AE, Murray GF, Perez V, Coulborn RM, Davis BM, Uddin M, et al. Mask use, hand hygiene, and seasonal influenza-like illness among young adults: a randomized intervention trial. *J Infect Dis*. 2010;201(4):491-8.
116. Aiello AE, Perez V, Coulborn RM, Davis BM, Uddin M, Monto AS. Facemasks, hand hygiene, and influenza among young adults: a randomized intervention trial. *PLoS One*. 2012;7(1):e29744.
117. World Health Organization. Information Note COVID-19 and NCDs. Geneva: World Health Organization. 2020. (https://www.who.int/docs/default-source/inaugural-who-partners-forum/covid-19-and-ncds---final---corr7.pdf?sfvrsn=9b65e287_1&download=true, accessed 21 November 2020)
118. World Health Organization. Coronavirus disease (COVID-19) advice for the public: When and how to use masks. Geneva: World Health Organization; 2020. (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public/when-and-how-to-use-masks>, accessed 21 November 2020).
119. Aydin O, Emon B, Cheng S, Hong L, Chamorro LP, Saif MTA. Performance of fabrics for home-made masks against the spread of COVID-19 through droplets: A quantitative mechanistic study. *Extreme Mech Lett*. 2020;40:100924.
120. Fischer EP, Fischer MC, Grass D, Henrion I, Warren WS, Westman E. Low-cost measurement of face mask efficacy for filtering expelled droplets during speech. *Sci Adv*. 2020;6(36).
121. Milton DK, Fabian MP, Cowling BJ, Grantham ML, McDevitt JJ. Influenza virus aerosols in human exhaled breath: particle size, culturability, and effect of surgical masks. *PLoS Pathog*. 2013;9(3):e1003205.
122. Bion JF, Abrusci T, Hibbert P. Human factors in the management of the critically ill patient. *Br J Anaesth*. 2010;105(1):26-33.
123. Chen YJ, Qin G, Chen J, Xu JL, Feng DY, Wu XY, et al. Comparison of Face-Touching Behaviors Before and During the Coronavirus Disease 2019 Pandemic. *JAMA Netw Open*. 2020;3(7):e2016924.
124. Shiraly R, Shayan Z, McLaws ML. Face touching in the time of COVID-19 in Shiraz, Iran. *Am J Infect Control*. 2020. 48(12): 1559–1561.
125. Betsch C, Korn L, Sprengholz P, Felgendreiff L, Eitze S, Schmid P, et al. Social and behavioral consequences of mask policies during the COVID-19 pandemic. *Proc Natl Acad Sci U S A*. 2020;117(36):21851-3.
126. Cowling BJ, Ali ST, Ng TWY, Tsang TK, Li JCM, Fong MW, et al. Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public Health*. 2020;5(5):e279-e88.
127. Giacalone S, Minuti A, Spigariolo CB, Passoni E, Nazzaro G. Facial dermatoses in the general population due to wearing of personal protective masks during the COVID-19 pandemic: first observations after lockdown. *Clin Exp Dermatol*. 2020. doi: 10.1111/ced.14376
128. Hufner K, Hofer A, Sperner-Unterweger B. On the difficulties of building therapeutic relationships when wearing face masks. *J Psychosom Res*. 2020;138:110226.
129. Crume B. The silence behind the mask: my journey as a deaf pediatric resident amidst a pandemic. *Acad Pediatr*. 2020. doi: 10.1016/j.acap.2020.10.002.
130. Allison MA, Guest-Warnick G, Nelson D, Pavia AT, Srivastava R, Gesteland PH, et al. Feasibility of elementary school children's use of hand gel and facemasks during influenza season. *Influenza Other Respir Viruses*. 2010;4(4):223-9.
131. Canini L, Andreoletti L, Ferrari P, D'Angelo R, Blanchon T, Lemaitre M, et al. Surgical mask to prevent influenza transmission in households: a cluster randomized trial. *PLoS One*. 2010;5(11):e13998.
132. Uchida M, Kaneko M, Hidaka Y, Yamamoto H, Honda T, Takeuchi S, et al. Effectiveness of vaccination and wearing masks on seasonal influenza in Matsumoto City, Japan, in the 2014/2015 season: An observational study among all elementary schoolchildren. *Prev Med Rep*. 2017;5:86-91.
133. Zand A, Heir A. Environmental impacts of new Coronavirus outbreak in Iran with an emphasis on waste management sector. *J Mater Cycles Waste Manag*. 2020 : 1–8.
134. Fikenzler S, Uhe T, Lavall D, Rudolph U, Falz R, Busse M, et al. Effects of surgical and FFP2/N95 face masks on cardiopulmonary exercise capacity. *Clin Res Cardiol*. 2020 Jul 6 : 1–9.
135. Harber P, Santiago S, Bansal S, Liu Y, Yun D, Wu S. Respirator physiologic impact in persons with mild respiratory disease. *J Occup Environ Med*. 2010;52(2):155-62.
136. Kyung SY, Kim Y, Hwang H, Park JW, Jeong SH. Risks of N95 Face Mask Use in Subjects With COPD. *Respir Care*. 2020;65(5):658-64.
137. Lee HP, Wang de Y. Objective assessment of increase in breathing resistance of N95 respirators on human subjects. *Ann Occup Hyg*. 2011;55(8):917-21.
138. Matuschek C, Moll F, Fangerau H, Fischer JC, Zanker K, van Griensven M, et al. Face masks: benefits and risks during the COVID-19 crisis. *Eur J Med Res*. 2020;25(1):32.
139. Person E, Lemerrier C, Royer A, Reychler G. [Effect of a surgical mask on six minute walking distance]. *Rev Mal Respir*. 2018;35(3):264-8.
140. Wong AY, Ling SK, Louie LH, Law GY, So RC, Lee DC, et al. Impact of the COVID-19 pandemic on sports and exercise. *Asia Pac J Sports Med Arthrosc Rehabil Technol*. 2020;22:39-44.
141. Li Y, Tokura H, Guo YP, Wong AS, Wong T, Chung J, et al. Effects of wearing N95 and surgical facemasks on heart rate, thermal stress and subjective sensations. *Int Arch Occup Environ Health*. 2005;78(6):501-9.
142. Hopkins SR, Dominelli PB, Davis CK, Guenette JA, Luks AM, Molgat-Seon Y, et al. Facemasks and the Cardiorespiratory Response to Physical Activity in Health and Disease. *Ann Am Thorac Soc*. 2020. doi:10.1513/AnnalsATS.202008-990CME.
143. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451-62.
144. Yang GZ, Kelley E, Darzi A. Patients' safety for global health. *Lancet*. 2011; 377(9769): 886-7.
145. Roberge RJ. Face shields for infection control: A review. *J Occup Environ Hyg*. 2016;13(4):235-42.

146. Jang JY, Kim, S.W. Evaluation of Filtration Performance Efficiency of Commercial Cloth Masks. *Journal of Environmental Health Sciences* (한국환경보건학회지)2015; 41 (3) 203-215.
147. Jung H, Kim JK, Lee S, Lee J, Kim J, Tsai P, et al. Comparison of Filtration Efficiency and Pressure Drop in Anti-Yellow Sand Masks, Quarantine Masks, Medical Masks, General Masks, and Handkerchiefs. *Aerosol Air Qual. Res.* 2014;14, 991–1002.
148. Lustig SR, Biswakarma JJH, Rana D, Tilford SH, Hu W, Su M, et al. Effectiveness of Common Fabrics to Block Aqueous Aerosols of Virus-like Nanoparticles. *ACS Nano.* 2020;14(6):7651-8.
149. Zangmeister CD, Radney JG, Vicenzi EP, Weaver JL. Filtration Efficiencies of Nanoscale Aerosol by Cloth Mask Materials Used to Slow the Spread of SARS-CoV-2. *ACS Nano.* 2020;14(7):9188-200.
150. Zhao M, Liao L, Xiao W, Yu X, Wang H, Wang Q, et al. Household materials selection for homemade cloth face coverings and their filtration efficiency enhancement with triboelectric charging. *Nano Lett.* 2020; 20(7):5544-5552.
151. Clase CM, Fu EL, Ashur A, Beale RCL, Clase IA, Dolovich MB, et al. Forgotten Technology in the COVID-19 Pandemic: Filtration Properties of Cloth and Cloth Masks-A Narrative Review. *Mayo Clin Proc.* 2020;95(10):2204-24.
152. Jain M, Kim S, Xu C, Li H, Rose G. Efficacy and Use of Cloth Masks: A Scoping Review. *Cureus* 12(9): e10423. doi:10.7759/cureus.10423
153. Mondal A, Das A, Goswami R. Utility of Cloth Masks in Preventing Respiratory Infections: A Systematic Review. *MedRxiv.* 2020 doi: 10.1101/2020.05.07.20093864
154. Roberge RJ, Roberge MR. Cloth face coverings for use as facemasks during the coronavirus (SARS-Cov-2) pandemic: what science and experience have taught us. *Disaster Med Public Health Prep.* 2020:1-29.
155. Sharma SK, Mishra M, Mudgal SK. Efficacy of cloth face mask in prevention of novel coronavirus infection transmission: A systematic review and meta-analysis. *J Educ Health Promot.* 2020;9:192.
156. Taminato M, Mizusaki-Imoto A, Saconato H, Franco E, Puga M, Duarte M, et al. Homemade cloth face masks as a barrier against respiratory droplets - systematic review. *Acta Paul Enferm.* 2020:eAPE20200103.
157. Bae S, Kim MC, Kim JY, Cha HH, Lim JS, Jung J, et al. Effectiveness of Surgical and Cotton Masks in Blocking SARS-CoV-2: A Controlled Comparison in 4 Patients. *Ann Intern Med.* 2020;173(1):W22-W3.
158. Ma QX, Shan H, Zhang HL, Li GM, Yang RM, Chen JM. Potential utilities of mask-wearing and instant hand hygiene for fighting SARS-CoV-2. *J Med Virol.* 2020. doi: 10.1002/jmv.25805.
159. Davies A, Thompson KA, Giri K, Kafatos G, Walker J, Bennett A. Testing the efficacy of homemade masks: would they protect in an influenza pandemic? *Disaster Med Public Health Prep.* 2013;7(4):413-8.
160. Konda A, Prakash A, Moss GA, Schmoldt M, Grant GD, Guha S. Aerosol Filtration Efficiency of Common Fabrics Used in Respiratory Cloth Masks. *ACS Nano.* 2020;14(5):6339-47.
161. Neupane BB, Mainali S, Sharma A, Giri B. Optical microscopic study of surface morphology and filtering efficiency of face masks. *PeerJ.* 2019;7:e7142.
162. Shakya KM, Noyes A, Kallin R, Peltier RE. Evaluating the efficacy of cloth facemasks in reducing particulate matter exposure. *J Expo Sci Environ Epidemiol.* 2017;27(3):352-7.
163. Jung H KJ, Lee S, Lee J, Kim J, Tsai P, et al. . Comparison of filtration efficiency and pressure drop in anti-yellow sand masks, quarantine masks, medical masks, general masks, and handkerchiefs. *Aerosol Air Qual Res.* 2014;14:991–1002.
164. Rengasamy S, Eimer B, Shaffer RE. Simple respiratory protection--evaluation of the filtration performance of cloth masks and common fabric materials against 20-1000 nm size particles. *Ann Occup Hyg.* 2010;54(7):789-98.
165. Dato VM, Hostler D, Hahn ME. Simple respiratory mask. *Emerg Infect Dis.* 2006;12(6):1033-4.
166. van der Sande M, Teunis P, Sabel R. Professional and home-made face masks reduce exposure to respiratory infections among the general population. *PLoS One.* 2008;3(7):e2618.
167. Chughtai AA, Seale H, Dung TC, Hayen A, Rahman B, Raina MacIntyre C. Compliance with the Use of Medical and Cloth Masks Among Healthcare Workers in Vietnam. *Ann Occup Hyg.* 2016;60(5):619-30.
168. AATCC. AATCC M14-2020 Guidance and Considerations for General Purpose Textile Face Coverings: Adult (<https://www.aatcc.org/covid/> accessed 28 November 2020)
169. Centers for Disease Control and Prevention. Scientific Brief: Community Use of Cloth Masks to Control the Spread of SARS-CoV-2. https://www.cdc.gov/coronavirus/2019-ncov/more/masking-science-sars-cov2.html?fbclid=IwAR28PppCa6x2uxwO8Z2baHM0KHS4JXx0inzMQs3zRHV1qq1_0a8mxZfpCw (Accessed 29 November 2020).
170. Swinfen R, Swinfen P. Low-cost telemedicine in the developing world. *J Telemed Telecare.* 2002;8(suppl 3):63-5.
171. Lee SA, Hwang DC, Li HY, Tsai CF, Chen CW, Chen JK. Particle Size-Selective Assessment of Protection of European Standard FFP Respirators and Surgical Masks against Particles-Tested with Human Subjects. *J Healthc Eng.* 2016;2016

Acknowledgments

This document was developed based on advice by the Strategic and Technical Advisory Group for Infectious Hazards (STAG-IH), and in consultation with the following members of:

- 1) The WHO Health Emergencies Programme (WHE) Ad-hoc COVID-19 IPC Guidance Development Group (in alphabetical order):

Jameela Alsaman, Ministry of Health, Bahrain; Anucha Apisarnthanarak, Thammasat University Hospital, Thailand; Baba Aye, Public Services International, France; Gregory Built, UNICEF, United States of America (USA); Roger Chou, Oregon Health Science University, USA; May Chu,

Colorado School of Public Health, USA; John Conly, Alberta Health Services, Canada; Barry Cookson, University College London, United Kingdom (U.K); Nizam Damani, Southern Health & Social Care Trust, United Kingdom; Dale Fisher, GOARN, Singapore; Joost Hopman, Radboud University Medical Center, The Netherlands; Mushtuq Husain, Institute of Epidemiology, Disease Control & Research, Bangladesh; Kushlani Jayatilleke, Sri Jayewardenapura General Hospital, Sri Lanka; Seto Wing Jong, School of Public Health, Hong Kong SAR, China; Souha Kanj, American University of Beirut Medical Center, Lebanon; Daniele Lantagne, Tufts University, USA; Fernanda Lessa, Centers for Disease Control and Prevention, USA; Anna Levin, University of São Paulo, Brazil; Ling Moi Lin, Sing Health, Singapore; Caline Mattar, World Health Professions Alliance, USA; Mary-Louise McLaws, University of New South Wales, Australia; Geeta Mehta, Journal of Patient Safety and Infection Control, India; Shaheen Mehtar, Infection Control Africa Network, South Africa; Ziad Memish, Ministry of Health, Saudi Arabia; Babacar Ndoeye, Infection Control Africa Network, Senegal; Fernando Otaiza, Ministry of Health, Chile; Diamantis Plachouras, European Centre for Disease Prevention and Control, Sweden; Maria Clara Padoveze, School of Nursing, University of São Paulo, Brazil; Mathias Pletz, Jena University, Germany; Marina Salvadori, Public Health Agency of Canada, Canada; Mitchell Schwaber, Ministry of Health, Israel; Nandini Shetty, Public Health England, United Kingdom; Mark Sobsey, University of North Carolina, USA; Paul Ananth Tambyah, National University Hospital, Singapore; Andreas Voss, Canisius-Wilhelmina Ziekenhuis, The Netherlands; Walter Zingg, University of Geneva Hospitals, Switzerland;

2) The WHO Technical Advisory Group of Experts on Personal Protective Equipment (TAG PPE):

Faisal Al Shehri, Saudi Food and Drug Authority, Saudi Arabi; Selcen Ayse, Istanbul University-Cerrahpasa, Turkey; Razan Asally, Saudi Food and Drug Authority, Saudi Arabi; Kelly Catlin, Clinton Health Access Initiative; Patricia Ching, WHO Collaborating Center, The University of Hong Kong, China; Mark Croes, Centexbel, Spring Gombe, United Nations; Emilio Hornsey, UK Public Health Rapid Support Team, U.K.; Selcen Kilinc-Balci, United States Centers for

Disease Control and Prevention (CDC), USA; Melissa Leavitt, Clinton Health Access Initiative; John McGhie, International Medical Corps; Claudio Meirovich, Meirovich Consulting; Mike Paddock, UNDP, Trish Perl, University of Texas Southwestern Medical Center, USA; Alain Prat, Global Fund, Ana Maria Rule, Johns Hopkins Bloomberg School of Public Health, U.S.A; Jitendar Sharma, Andra Pradesh MedTech Zone, India; Alison Syrett, SIGMA, Reiner Voelksen, VOELKSEN Regulatory Affairs, Nasri Yussuf, IPC Kenya.

3) External IPC peer review group:

Paul Hunter, University of East Anglia, U.K; Direk Limmathurotsakul, Mahidol University, Thailand; Mark Loeb, Department of Pathology and Molecular Medicine, McMaster University, Canada; Kalisavar Marimuthu, National Centre for Infectious Diseases, Singapore; Yong Loo Lin School of Medicine, National University of Singapore; Nandi Siegfried, South African Medical Research Council, South Africa.

4) UNICEF observers: Nagwa Hasanin, Sarah Karmin, Raoul Kamadjeu, Jerome Pfaffmann,

WHO Secretariat:

Benedetta Allegranzi, Gertrude Avortri, Mekdim Ayana, Hanan Balkhy, April Baller, Elizabeth Barrera-Cancedda, Anjana Bhushan, Whitney Blanco, Sylvie Briand, Alessandro Cassini, Giorgio Cometto, Ana Paula Coutinho Rehse, Carmem Da Silva, Nino Dal Dayanguirang, Sophie Harriet Dennis, Sergey Eremin, Luca Fontana, Dennis Falzon, Nathan Ford, Nina Gobat, Jonas Gonseth-Garcia, Rebeca Grant, Tom Grein, Ivan Ivanov, Landry Kabego, Catherine Kane, Pierre Claver Kariyo, Ying Ling Lin, Ornella Lincetto, Abdi Mahamud, Madison Moon, Takeshi Nishijima, Kevin Babila Ousman, Pillar Ramon-Pardo, Paul Rogers, Nahoko Shindo, Alice Simniceanu, Valeska Stempliuk, Maha Talaat Ismail, Joao Paulo Toledo, Anthony Twywan, Maria Van Kerkhove, Adriana Velazquez, Vicky Willet, Masahiro Zakoji, Bassim Zayed.

WHO continues to monitor the situation closely for any changes that may affect this interim guidance. Should any factors change, WHO will issue a further update. Otherwise, this interim guidance document will expire 1 year after the date of publication.

Annex: Updated guidance on non-medical (fabric) masks

Background

A non-medical mask, also called fabric mask, community mask or face covering, is neither a medical device nor personal protective equipment. Non-medical masks are aimed at the general population, primarily for protecting others from exhaled virus-containing droplets emitted by the mask wearer. They are not regulated by local health authorities or occupational health associations, nor is it required for manufacturers to comply with guidelines established by standards organizations. Non-medical masks may be homemade or manufactured. The essential performance parameters include good breathability, filtration of droplets originating from the wearer, and a snug fit covering the nose and mouth. Exhalation valves on masks are discouraged as they bypass the filtration function of the mask.

Non-medical masks are made from a variety of woven and non-woven fabrics, such as woven cotton, cotton/synthetic blends, polyesters and breathable spunbond polypropylene, for example. They may be made of different combinations of fabrics, layering sequences and available in diverse shapes. Currently, more is known about common household fabrics and combinations to make non-medical masks with target filtration efficiency and breathability (119, 146-150). Few of these fabrics and combinations have been systematically evaluated and there is no single design, choice of material, layering or shape among available non-medical masks that are considered optimal. While studies have focussed on single fabrics and combinations, few have looked at the shape and universal fit to the wearer. The unlimited combination of available fabrics and materials results in variable filtration and breathability.

In the context of the global shortage of medical masks and PPE, encouraging the public to create their own fabric masks may promote individual enterprise and community integration. Moreover, the production of non-medical masks may offer a source of income for those able to manufacture masks within their communities. Fabric masks can also be a form of cultural expression, encouraging public acceptance of protection measures in general. The safe re-use of fabric masks will also reduce costs and waste and contribute to sustainability (151-156).

This Annex is destined intended for two types of readers: homemade mask makers and factory-made masks manufacturers. Decision makers and managers (national/sub-national level) advising on a type of non-medical mask are also the focus of this guidance and should take into consideration the following features of non-medical masks: breathability, filtration efficiency (FE), or filtration, number and combination of fabric layers material used, shape, coating and maintenance.

Evidence on the effectiveness of non-medical (fabric) masks

A number of reviews have been identified on the effectiveness of non-medical masks (151-156). One systematic review (155) identified 12 studies and evaluated study quality. Ten were laboratory studies (157-166), and two reports were from a single randomized trial (72, 167). The majority of studies were conducted before COVID-19 emerged or used laboratory generated particles to assess filtration efficacy. Overall, the reviews concluded that

cloth face masks have limited efficacy in combating viral infection transmission.

Homemade non-medical masks

Homemade non-medical masks made of household fabrics (e.g., cotton, cotton blends and polyesters) should ideally have a three-layer structure, with each layer providing a function (see Figure 1) (168). It should include:

1. an innermost layer (that will be in contact with the face) of a hydrophilic material (e.g., cotton or cotton blends of terry cloth towel, quilting cotton and flannel) that is non-irritating against the skin and can contain droplets (148)
2. a middle hydrophobic layer of synthetic breathable non-woven material (spunbond polypropylene, polyester and polyaramid), which may enhance filtration, prevent permeation of droplets or retain droplets (148, 150)
3. an outermost layer made of hydrophobic material (e.g. spunbond polypropylene, polyester or their blends), which may limit external contamination from penetrating through the layers to the wearer's nose and mouth and maintains and prevents water accumulation from blocking the pores of the fabric (148).

Although a minimum of three layers is recommended for non-medical masks for the most common fabric used, single, double or other layer combinations of advanced materials may be used if they meet performance requirements. It is important to note that with more tightly woven materials, breathability may be reduced as the number of layers increases. A quick check may be performed by attempting to breathe, through the mouth, through the multiple layers.



Figure 1. Non-medical mask construction using breathable fabrics such as cotton, cotton blends, polyesters, nylon and polypropylene spunbond that are breathable may impart adequate filtration performance when layered. Single- or double-layer combinations of advanced materials may be used if they meet performance requirements (72).

Assumptions regarding homemade masks are that individual makers only have access to common household fabrics and do not have access to test equipment to confirm target performance (filtration and breathability). Figure 1 illustrates a multi-layer mask construction with examples of fabric options. Very porous materials, such as gauze, even with multiple layers, may provide very low filtration efficiency (147). Higher thread count fabrics offer improved filtration performance (169). Coffee filters, vacuum bags and materials not meant for clothing should be avoided as they may contain injurious content when breathed in. Microporous films such as Gore-Tex are not recommended (170).

Factory-made non-medical masks: general considerations for manufacturers

The non-medical mask, including all components and packaging, must be non-hazardous, non-toxic and child-friendly (no exposed sharp edges, protruding hardware or rough materials). Factory-made non-medical masks must be made using a process that is certified to a quality management system (e.g., ISO 9001). Social accountability standards (e.g., SAI SA8000) for multiple aspects of fair labour practices, health and safety of the work force and adherence to UNICEF's Children's Rights and Business Principles are strongly encouraged.

Standards organizations' performance criteria

Manufacturers producing masks with consistent standardized performance can adhere to published, freely available guidance from several organizations including those from: the French Standardization Association (AFNOR Group), The European Committee for Standardization (CEN), Swiss National COVID-19 Task Force, the American Association of Textile Chemists and Colorists (AATCC), the South Korean Ministry of Food and Drug Safety (MFDS), the Italian Standardization Body (UNI) and the Government of Bangladesh.

Essential parameters

The essential parameters presented in this section are the synthesis of the abovementioned regional and national guidance. They include filtration, breathability and fit. Good performance is achieved when the three essential parameters are optimized at the preferred threshold (Figure 2).




Figure 2. Illustration of the three essential parameters of filtration, breathability and fit.

The summary of the three essential parameters can be found in Table 1 and the additional performance considerations in Table 2. The minimum threshold is the minimum acceptable parameter, while the preferred threshold is the optimum.

Filtration and breathability

Filtration depends on the filtration efficiency (in %), the type of challenge particle (oils, solids, droplets containing bacteria) and the particle size (see Table 1). Depending on the fabrics used, filtration and breathability can complement or work against one another. The selection of material for droplet filtration (barrier) is as important as breathability. Filtration is dependent on the tightness of the weave, fibre or thread diameter. Non-woven materials used for disposable masks are manufactured using processes to create polymer fibres that are thinner than natural fibres such as cotton and that are held together by partial melting.

Breathability is the difference in pressure across the mask and is typically reported in millibars (mbar) or Pascals (Pa) or, normalized to the cm^2 in mbar/cm^2 or Pa/cm^2 . Acceptable breathability of a medical mask should be below $49 \text{ Pa}/\text{cm}^2$. For non-medical masks, an acceptable pressure difference, over the whole mask, should be below $60 \text{ Pa}/\text{cm}^2$, with lower values indicating better breathability.

Non-medical fabric masks consisting of two layers of polypropylene spunbond and two layers of cotton have been shown to meet the minimum requirements for droplet filtration and breathability of the CEN CWA 17553 guidance. It is preferable not to select elastic material to make masks as the mask material may be stretched over the face, resulting in increased pore size and lower filtration through multiple usage. Additionally, elastic fabrics are sensitive to washing at high temperatures thus may degrade over time.

Coating the fabric with compounds like wax may increase the barrier and render the mask fluid resistant; however, such coatings may inadvertently completely block the pores and make the mask difficult to breathe through. In addition to decreased breathability unfiltered air may more likely escape the sides of the mask on exhalation. Coating is therefore not recommended.

Valves that let unfiltered air escape the mask are discouraged and are an inappropriate feature for masks used for the purpose of preventing transmission.

Table 1. Essential parameters (minimum and preferred thresholds) for manufactured non-medical mask

Essential Parameters	Minimum threshold	Preferred threshold
1. Filtration*		
1.1. filtration efficiency	70% @ 3 micron	> 70%, without compromising breathability
1.2. Challenge particle	Solid: sodium chloride (NaCl), Talcum powder, Holi powder, dolomite, Polystyrene Latex spheres Liquid: DEHS Di-Ethyl-Hexyl-Sebacat, paraffin oil	Based on availability
1.3. Particle size	Choose either sizes: 3 µm, 1 µm, or smaller	Range of particle sizes
2. Breathability		
2.1. Breathing resistance**	≤60 Pa/cm ²	Adult: ≤ 40 Pa/cm ² Paediatric: ≤ 20 Pa/cm ²
2.2 Exhalation valves	Not recommended	N/A
3. Fit		
3.1. Coverage	Full coverage of nose and mouth, consistent, snug perimeter fit at the nose bridge, cheeks, chin and lateral sides of the face; adequate surface area to minimize breathing resistance and minimize side leakage	Same as current requirements
3.2 Face seal	Not currently required	Seal as good as FFR (respirator): Fit factor of 100 for N95 Maximum Total Inward Leakage of 25% (FFP1 requirement)
3.2. Sizing	Adult and child	Should cover from the bridge of the nose to below the chin and cheeks on either side of the mouth Sizing for adults and children (3-5, 6-9, 10-12, >12)
3.3 Strap strength		> 44.5 N

* Smaller particle may result in lower filtration.

** High resistance can cause bypass of the mask. Unfiltered air will leak out the sides or around the nose if that is the easier path.

Fit: shape and sizing

Fit is the third essential parameter, and takes into consideration coverage, seal, sizing, and strap strength. Fit of masks currently is not defined by any standard except for the anthropometric considerations of facial dimensions (ISO/TS 16976-2) or simplified to height mask (South Korean standard for KF-AD). It is important to ensure that the mask can be held in place comfortably with as little adjustment of the elastic bands or ties as possible.

Mask shapes typically include flat-fold or duckbill and are designed to fit closely over the nose, cheeks and chin of the wearer. Snug fitting designs are suggested as they limit leaks of unfiltered air escaping from the mask (148). Ideally the mask should not have contact with the lips, unless hydrophobic fabrics are used in at least one layer of the mask (148). Leaks where unfiltered air moves in and out of the mask may be attributed to the size and shape of the mask (171).

Additional considerations

Optional parameters to consider in addition to the essential performance parameters include if reusable, biodegradability for disposal masks, antimicrobial performance where applicable and chemical safety (see Table 2).

Non-medical masks intended to be reusable should include instructions for washing and must be washed a minimum of five cycles, implying initial performance is maintained after each wash cycle.

Advanced fabrics may be biodegradable or compostable at the end of service life, according to a recognized standard process (e.g., UNI EN 13432, UNI EN 14995 and UNI / PdR 79).

Manufacturers sometimes claim their NM masks have antimicrobial performance. Antimicrobial performance may be due to coatings or additives to the fabric fibres. Treated fabrics must not come into direct contact with mucous membranes; the innermost fabric should not be treated with

antimicrobial additives, only the outermost layer. In addition, antimicrobial fabric standards (e.g., ISO 18184, ISO 20743, AATCC TM100, AATCC 100) are generally slow acting. The inhibition on microbial growth may take full effect after 2- or 24-hour contact time depending on the standard. The standards have generally been used for athletic apparel and substantiate claims of odour control performance. These standards are not appropriate for non-medical cloth masks and may provide a false sense of protection from infectious agents. If claims are made, manufacturers should specify which standard supports antimicrobial performance, the challenge organism and the contact time.

Volatile additives are discouraged as these may pose a health risk when inhaled repeatedly during wear. Certification according to organizations including OEKO-TEX (Europe) or SEK (Japan), and additives complying with REACH (Europe) or the Environmental Protection Agency (EPA, United States of America) indicate that textile additives are safe and added at safe levels.

Table 2. Additional parameters for manufactured non-medical masks

Additional parameters	Minimum thresholds
If reusable, number of wash cycles	5 cycles
Disposal	Reusable If biodegradable (CFC-BIO), according to UNI EN 13432, UNI EN 14995
Antimicrobial (bacteria, virus, fungus) performance	ISO 18184 (virus) ISO 20743 (bacteria) ISO 13629 (fungus) AATCC TM100 (bacteria)
Chemical safety	Comply with REACH regulation, including inhalation safety